

Feed Mix Baby Calf Pellet

Medicated

Supplemental Feed for Baby Calves

Active Drug Ingredient

Decoquinatone.....54.4 g/ton
Chlortetracycline.....50g/ton

Guaranteed Analysis

| | | | |
|---------------|---------------|--------|-------|
| Crude Protein | not less than | 14.0 | % |
| Crude Fat | not less than | 2.5 | % |
| Crude Fiber | not more than | 10.0 | % |
| Calcium | not less than | 0.5 | % |
| Calcium | not more than | 0.9 | % |
| Phosphorus | not less than | 0.4 | % |
| Salt | not less than | 0.1 | % |
| Salt | not more than | 0.5 | % |
| Potassium | not less than | 0.6 | % |
| Vitamin A | not less than | 3600.0 | IU/lb |
| Vitamin D | not less than | 360.0 | IU/lb |
| Vitamin E | not less than | 12.0 | IU/lb |
| Magnesium | not less than | 0.2 | % |
| Copper | not less than | 16.0 | ppm |
| Selenium | not less than | 0.40 | ppm |
| Zinc | not less than | 80.0 | ppm |

Ingredients

Processed Grain By-Products, Grain Products, Calcium Carbonate, Potassium Chloride, Salt, Magnesium Oxide, Vegetable Fat, Manganese Hydroxychloride, Zinc Hydroxychloride, Selenium Yeast, Basic Copper Chloride, Vitamin E Supplement, Ethylenediamine Dihydriodide, Cobalt Proteinate, Thiamin Hydrochloride, Vitamin A Supplement, Vitamin D Supplement, Yea-Sacc 1026®, Bio-Mos®, Molasses Flavoring

Directions for Use

Feed Mix Baby Calf Pellet is designed for young calves. It contains the right amount of protein, energy, vitamins, minerals, yeast culture and antibiotics to maintain digestive and immune health. Feed Mix Baby Calf Pellet is a supplemental feed to mother's milk or milk replacer.

Begin feeding Feed Mix Baby Calf Pellet at 4 days of age and continue until calves are 8 to 12 weeks old. At this time they may be switched to Feed Mix Creep Feed or Feed Mix Creep Pellet.

Good quality forage should be available. Fresh, clean water should always be provided.

Feed Mix Baby Calf Feed can be fed to prevent coccidiosis in ruminating and non-ruminating calves (including veal calves) and cattle caused by Eimeria bovis and E. zuernii. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Coccidostats are not indicated for use in adult animals due to continuous previous exposure.

Caution

Do not feed to sheep due to high copper levels. Do not feed to cows providing milk for food. A withdrawal period for Chlortetracycline has not been established in pre-ruminating calves. Do not use Chlortetracycline in calves to be processed for veal. Do not feed Chlortetracycline to ducks or turkeys producing eggs for human consumption.

Manufactured by:



HARVEY, ND

414 7th St W

Harvey, ND 58341

701-324-2201

Net Weight on Invoice

This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.