

Feed Mix Finish Feed Medicated

Supplemental Feed for Cattle at Finishing
Weights

Active Drug Ingredient

Monensin.....36g/ton

Guaranteed Analysis

Crude Protein	not less than	13.0	%
Crude Fat	not less than	4.0	%
Crude Fiber	not more than	11.0	%
Calcium	not less than	0.5	%
Calcium	not more than	0.9	%
Phosphorus	not less than	0.4	%
Salt	not less than	0.1	%
Salt	not more than	0.5	%
Potassium	not less than	0.8	%
Vitamin A	not less than	3600.0	IU/lb
Vitamin D	not less than	360.0	IU/lb
Vitamin E	not less than	20.0	IU/lb
Magnesium	not less than	0.2	%
Copper	not less than	19.0	ppm
Selenium	not less than	0.40	ppm
Zinc	not less than	90.0	ppm

Ingredients

Grain Products, Processed Grain By-Products, Calcium Carbonate, Potassium Chloride, Salt, Magnesium Oxide, Vegetable Fat, Manganese Hydroxychloride, Zinc Hydroxychloride, Selenium Yeast, Basic Copper Chloride, Vitamin E Supplement, Ethylenediamine Dihydriodide, Cobalt Proteinate, Thiamin Hydrochloride, Vitamin A Supplement, Vitamin D Supplement, Yea-Sacc 1026 ®

Directions for Use

Feed Mix Finishing Feed is designed for cattle being finished for slaughter. All ration changes should be made gradually to prevent digestive disturbances. Feed Mix Finishing Feed at a rate of 2% of body weight. Unlimited roughage in the form of hay or grass should be available, or add hay at 1% body weight for a TMR. Fresh, clean water should be available at all times.

To provide 50 to 200 milligrams of Monensin feed at a rate of 2.3 to 8.9 pounds daily.

Caution

Do not feed to sheep due to high copper levels. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured by:



Feed Mix
414 7th St W
Harvey, ND 58341
701-324-2201

Net Weight on Invoice

This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.