

Feed Mix Creep Pellet Medicated

Supplemental Feed for Young Calves on Pasture

Active Drug Ingredient

Monensin..... 45 g/ton

Guaranteed Analysis

Crude Protein	not less than	14.0	%
Crude Fat	not less than	3.0	%
Crude Fiber	not more than	10.0	%
Calcium	not less than	0.5	%
Calcium	not more than	0.9	%
Phosphorus	not less than	0.5	%
Salt	not less than	0.1	%
Salt	not more than	0.5	%
Potassium	not less than	0.7	%
Vitamin A	not less than	3600.0	IU/lb
Vitamin D	not less than	360.0	IU/lb
Vitamin E	not less than	12.0	IU/lb
Magnesium	not less than	0.2	%
Copper	not less than	16.0	ppm
Selenium	not less than	0.50	ppm
Zinc	not less than	88.0	ppm

Ingredients

Processed Grain By-Products, Grain Products, Molasses Products, Lignin Sulfonate, Calcium Carbonate, Potassium Chloride, Salt, Magnesium Oxide, Vegetable Fat, Manganese Hydroxychloride, Zinc Hydroxychloride, Selenium Yeast, Basic Copper Chloride, Vitamin E Supplement, Ethylenediamine Dihydriodide, Cobalt Proteinate, Thiamin Hydrochloride, Vitamin A Supplement, Vitamin D Supplement, Yea-Sacc 1026®, Molasses Flavoring

Directions for Use

Feed Mix Creep Pellet is a supplement for calves on pasture or range grasses. Providing a supplement feed while on pasture may result in increased weights and easier transitions to eating grains at weaning.

Feed Mix Creep Pellet may be provided from 10 days of age until calves are weaned.

Unlimited roughage in the form of hay or pasture should be provided.

Fresh, clean water should be available at all times.

Feed FM Creep Pellet continuously at a rate of 2.3 to 8.9 pounds daily to provide 50 to 200 milligrams of Monensin.

Caution

Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to sheep due to high copper levels.

Manufactured by:

FEED MIX

PRECISION LIVESTOCK NUTRITION

HARVEY, ND

414 7th St W

Harvey, ND 58341

701-324-2201

Net Weight on Invoice

This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.