Feed Mix Weaning Feed Medicated

Supplemental Feed for Newly Weaned Calves

Active Drug Ingredient

Lasalocid	45 g/ton
Chlortetracycline	125 g/ton

	Cuaranteeu / marysis			
Crude Protein	not less than	14.0	%	
Crude Fat	not less than	4.0	%	
Crude Fiber	not more than	11.0	%	
Calcium	not less than	0.5	%	
Calcium	not more than	0.9	%	
Phosphorus	not less than	0.5	%	
Salt	not less than	0.1	%	
Salt	not more than	0.5	%	
Potassium	not less than	0.9	%	
Vitamin A	not less than	3600.0	IU/lb	
Vitamin D	not less than	360.0	IU/lb	
Vitamin E	not less than	39.0	IU/lb	
Magnesium	not less than	0.3	%	
Copper	not less than	20.0	ppm	
Selenium	not less than	0.60	ppm	
Zinc	not less than	104.0	ppm	

Guaranteed Analysis

Ingredients

Processed Grain By-Products, Grain Products, Calcium Carbonate, Potassium Chloride, Salt, Magnesium Oxide, Vegetable Fat, Manganese Hydroxychloride, Zinc Hydroxychloride, Selenium Yeast, Basic Copper Chloride, Vitamin E Supplement, Ethylenediamine Dihydriodide, Cobalt Proteinate, Thiamin Hydrochloride, Vitamin A Supplement, Vitamin D Supplement, Bio-Mos®, Yea-Sacc 1026®

Directions for Use

Feed Mix Weaning Feed is designed as a supplemental feed for newly weaned calves. Feed Mix Weaning Feed helps support digestive and immune health of stressed calves by having proper levels of energy, protein, vitamins, and minerals. Feed at a rate of 2% of body weight. Hay should be fed at a rate of 1% of body weight. Fresh clean water should be available at all times. To provide 100 to 360 milligrams Lasalocid feed FM Weaning Feed at a rate of 4.5 to 16 pounds daily. To provide 360 milligrams of Chlortetracycline feed FM Weaning Feed at a rate of 16 pounds per head per day.

Caution

Do not allow horses or other equine access to premixes or supplements containing lasalocid, as ingestion may be fatal. The safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors resulting in excessive concentrations of lasalocid could be fatal to cattle and sheep. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for yeal. Do not feed to sheep due to high copper levels. A withdrawal period has not been established for this product in pre-ruminant calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.

Manufactured By:



414 7th St W Harvey, ND 58341 701-324-2201

Net Weight on Invoice

This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.